

MERS-CoV Spike Protein S1 Antibody (C-terminal), Rabbit PAb, Antigen Affinity Purified

Catalog_no :	AD-PD400062
Applications :	WB
Category :	冠状病毒产品
Size :	50μL/100μL
Immunogen :	A synthetic peptide corresponding to the C-terminus of the S1 subunit of MERS-CoV (NCoV / Novel coronavirus) spike glycoprotein (S protein).
Specificity :	MERS-CoV CoV spike
Source :	A synthetic peptide corresponding to the C-terminus of the S1 subunit of MERS-CoV (NCoV / Novel coronavirus) spike glycoprotein (S protein).
Dilution :	WB : 1:500-1:2000
Purification :	Protein A & Antigen Affinity
Background :	<p>The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that 2019-nCoV can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.</p>
缓冲液 :	0.2 μm filtered solution in PBS
运输及保存条件 :	This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
classification_1 :	Anti-coronavirus s1 Antibody; Anti-coronavirus s2 Antibody; Anti-coronavirus spike Antibody; Anti-cov spike Antibody; Anti-ncov RBD Antibody; Anti-ncov s1 Antibody; Anti-ncov s2 Antibody; Anti-ncov spike Antibody; Anti-RBD Antibody; Anti-S Antibody;



storage :	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
偶联物 :	Unconjugated
状态 :	Liquid