




MUC5AC (9X2) mouse mAb

| | |
|---------------------|---|
| Catalog_no : | AM6198 |
| Applications : | IHC-p |
| Reactivity : | Human |
| Category : | 抗原抗体 |
| Size : | 50µl/100µl |
| Gene_name : | MUC5AC MUC5 |
| Protein_name : | Mucin-5AC (MUC-5AC) (Gastric mucin) (Lewis B blood group antigen) (LeB) (Major airway glycoprotein) (Mucin-5 subtype AC, tracheobronchial) (Tracheobronchial mucin) (TBM) (Fragments) |
| Humanswissprot_no : | P98088 |
| Immunogen : | Synthesized peptide derived from human MUC5AC |
| Specificity : | This antibody detects endogenous levels of human MUC5AC |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Mouse/IgG1, Kappa |
| Dilution : | IHC-p 1:100-500 |
| Purification : | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |
| Notes : | apoptosis, induction of apoptosis, cell adhesion, digestion, cell death, negative regulation of cell proliferation,regulation of cell death, positive regulation of cell death, programmed cell death, induction of programmed cell death,death, biological |

Product Images



Immunohistochemical analysis of paraffin-embedded Stomach. 1, Antibody was diluted at 1:200(4°,overnight). 2, Tris-EDTA,pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).