




Prostatic Acid Phosphatase(PSAP) (8U8) mouse mAb

| | |
|---------------------|--|
| Catalog_no : | AM6213 |
| Applications : | IHC-p |
| Reactivity : | Human |
| Category : | 抗原抗体 |
| Size : | 50μl/100μl |
| Gene_name : | ACPP |
| Protein_name : | Prostatic acid phosphatase (PAP) (EC 3.1.3.2) (5'-nucleotidase) (5'-NT) (EC 3.1.3.5) (Ecto-5'-nucleotidase) (Thiamine monophosphatase) (TMPase) [Cleaved into: PAPf39] |
| Humanswissprot_no : | P15309 |
| Immunogen : | Synthesized peptide derived from human Prostatic Acid Phosphatase(PSAP) |
| Specificity : | This antibody detects endogenous levels of human Prostatic Acid Phosphatase(PSAP) |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Mouse/IgG1, Kappa |
| Dilution : | IHC-p 1:100-500 |
| Purification : | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage_stability : | -20°C/1 year |

Product Images



Immunohistochemical analysis of paraffin-embedded Benign Prostatic Hyperplasia. 1, Antibody was diluted at 1:200(4°,overnight). 2, Tris-EDTA,pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).